arXiv:1206.0382v1 [math.MG] 2 Jun 2012

BOUNDARIES OF DISK-LIKE SELF-AFFINE TILES

KING-SHUN LEUNG AND JUN JASON LUO

Abstract. Let $T := T(A, \mathcal{D})$ be a disk-like self-affine tile generated by an integral expanding matrix A and a consecutive collinear digit set $\mathcal{D}$, and let $f(x) = x^2 + px + q$ be the characteristic polynomial of A. By investigating the neighbors of T, we construct a neighbor graph and show that any point on the boundary ∂T can be identified with a one-sided infinite path of the graph. Also we derive certain equivalent conditions for the pair $(A, \mathcal{D})$ to be a number system. From the neighbor graph we can deduce a family of graph-directed sets, which gives a second description of ∂T. When A is a similarity, we obtain $\dim_H(\partial T) = 2\log\rho/\log|q|$ where ρ is the largest positive zero of the cubic polynomial $x^3 - (|p|-1)x^2 - (|q|-|p|)x - |q|$.

1. **Introduction**

Let $M_n(\mathbb{Z})$ denote the set of $n \times n$ matrices with entries in $\mathbb{Z}$ and let $A \in M_n(\mathbb{Z})$ be expanding (i.e., all eigenvalues of A have moduli > 1). Assume $|\det(A)| = |q|$, and $\mathcal{D} = \{0, d_1, \ldots, d_{|q|-1}\} \subset \mathbb{Z}^n$ with $|q|$ distinct vectors. We call $\mathcal{D}$ a *digit set* and $(A, \mathcal{D})$ a *self-affine pair*. It is well-known that there exists a unique self-affine set $T := T(A, \mathcal{D})$ [LW1] satisfying

$$T = A^{-1}(T + \mathcal{D}) = \left\{ \sum_{i=1}^{\infty} A^{-i} d_{j_i} : d_{j_i} \in \mathcal{D} \right\}.$$

If T has non-void interior, then there exists a subset $\mathcal{J} \subset \mathbb{Z}^n$ such that

$$T + \mathcal{J} = \mathbb{R}^n \quad \text{and} \quad (T+t)^\circ \cap (T+t')^\circ = \emptyset,\ t \neq t',\ t, t' \in \mathcal{J},$$

thus T is called a *self-affine tile* and $\mathcal{J}$ a tiling set. $T + \mathcal{J}$ is called a tiling of $\mathbb{R}^n$, and a lattice tiling if $\mathcal{J}$ is a lattice [LW3].

The topological properties of self-affine tiles, such as connectedness or disk-likeness (i.e., homeomorphic to the closed unit disk), have attracted a lot of interest. A systematical study on connectedness was due to Kirat and Lau [KL], they mainly concerned a class of self-affine tiles T generated by the *consecutive collinear (CC)* digit sets $\mathcal{D} := \mathcal{D}(v, |q|) = \{0, 1, \ldots, |q|-1\}v, v \in \mathbb{Z}^n \setminus \{0\}$ via the algebraic property of the characteristic polynomial of the matrix A. Other considerations on non-consecutive collinear or non-linear digit sets can be found in [DL], [LLu]. The

---

Date: March 6, 2013.

Key words and phrases. boundary, self-affine tile, disk-like, CC digit set, neighbor graph, contact matrix, graph-directed sets.

question of disk-likeness was first investigated by Bandt and Gelbrich [BG] for self-affine tiles in $\mathbb{R}^2$ with $|\det(A)| = 2$ or 3. They also observed that the characteristic polynomial of an expanding matrix $A \in M_2(\mathbb{Z})$ is of the form:

$$f(x) = x^2 + px + q, \text{ with } |p| \le q, \text{ if } q \ge 2; \quad |p| \le |q+2|, \text{ if } q \le -2.$$

By studying the neighborhood structure of T, Bandt and Wang [BW] proved that a tile T with no more than six neighbors is disk-like if and only if T is connected. A translation of the tile $T+\ell$, $\ell \in \mathcal{J}$ is called a *neighbor* of T if $T \cap (T+\ell) \neq \emptyset$. Making use of this criterion, Leung and Lau [LL] then gave a complete characterization of the disk-likeness of self-affine tiles with CC digit sets.

Theorem 1.1. *([LL]) Let $A \in M_2(\mathbb{Z})$ be an expanding matrix with characteristic polynomial $f(x) = x^2 + px + q$. Then for any CC digit set $\mathcal{D}(v, |q|)$ in $\mathbb{Z}^2$ such that v, Av are linearly independent, T is a disk-like tile if and only if $2|p| \le |q+2|$.*

Moreover, when $p = 0$, T is a square tile; when $p \neq 0$, T is a hexagonal tile.

Recently, Akiyama and Loridant ([AL], [AL2]) proposed a method to parameterize the boundary of a self-affine tile, and reproved Theorem 1.1 by showing that the boundary of T is a simple closed curve. In the present paper, we go further to explore the structure of the boundary of the T defined in Theorem 1.1. For convenience, we call such T a *CC tile*. If it is also disk-like, we call it a *disk-like CC tile*.

As the first main result of this paper, we describe the boundary ∂T of a disk-like CC tile T in two equivalent ways. First we define a labeled directed graph (called the neighbor graph of T) with $\mathcal{V} = \{\ell : T + \ell \text{ is a neighbor of } T\}$ as the set of vertices and $\Delta\mathcal{D}$ the set of edge labels, such that ∂T is identified as the union of all one-sided infinite paths of this graph. Hence ∂T determines a sofic system [Fi]. The neighbor graph is associated with a contact matrix M whose entries are the edge labels. Then we provide some equivalent conditions for the self-affine pair $(A, \mathcal{D})$ to be a number system [MTT]. On the other hand, ∂T can also be expressed as a family of graph-directed sets [SW] by using an algorithm (Proposition 3.1). We prove that the associated graph-directed IFS satisfies the open set condition. From this we can deduce our second main result, an expression for the Hausdorff dimension of ∂T. The derivation involves finding the Perron-Frobenius eigenvalue of M. When A is a similarity, we obtain the following.

Theorem 1.2. *Let $A \in M_2(\mathbb{Z})$ be an expanding similarity with characteristic polynomial $f(x) = x^2 + px + q$ and $T = T(A, \mathcal{D})$ be a disk-like CC tile. Then*

$$\dim_H(\partial T) = \frac{2\log\rho}{\log|q|}$$

where ρ is the largest positive zero of the cubic polynomial $x^3 - (|p|-1)x^2 - (|q| - |p|)x - |q|$.

The rest of the paper is organized as follows: in Section 2, we construct a neighbor graph and identify ∂T with a sofic system determined by this graph, moreover we derive some equivalent conditions for the self-affine pair $(A, \mathcal{D})$ to be a number system; in Section 3, we consider ∂T as graph-directed sets and prove Theorem 1.2 by checking the spectral radius of the contact matrix. Finally all neighbor graphs, graph-directed sets and contact matrices corresponding to different characteristic polynomials $f(x)$ are listed in Appendices A, B and C for easy reference.

2. **Boundary of T as a Sofic System**

We first introduce some terminology of symbolic dynamics from [LM]. Let $\mathcal{G} = \mathcal{G}(\mathcal{V}, \mathcal{E})$ be a directed graph where $\mathcal{V}$ is the set of vertices and $\mathcal{E}$ the set of edges. Let $\mathcal{A}$ be a finite set (called *alphabet*). If there exists a mapping (called *labeling*) $\mathcal{L} : \mathcal{E} \to \mathcal{A}$, then the ordered pair $\mathbf{G} = (\mathcal{G}, \mathcal{L})$ is called a *labeled directed graph.* All the infinite paths $\xi = e_1e_2e_3\ldots$ on $\mathcal{G}$ constitute the so-called *edge shift* $\mathbf{X}_{\mathcal{G}}$. Define the *label of the path* ξ by

$$\mathcal{L}_\infty(\xi) := \mathcal{L}(e_1)\mathcal{L}(e_2)\mathcal{L}(e_3)\ldots \in \mathcal{A}^{\mathbb{N}}.$$

Here $\mathcal{A}^{\mathbb{N}}$ is called the *full shift* of $\mathcal{A}$. The set of all such labels is denoted by

$$\mathbf{X}_{\mathbf{G}} = \left\{x \in \mathcal{A}^{\mathbb{N}} : x = \mathcal{L}_\infty(\xi) \text{ for some } \xi \in \mathbf{X}_{\mathcal{G}}\right\}.$$

Any subset of $\mathcal{A}^{\mathbb{N}}$ which can be defined by a labeled directed graph as above, is called a *sofic shift* or *sofic system* ([Fi], [LM]). Weiss [We] coined the term *sofic* which is derived from the Hebrew word for *finite* [LM].

Let $D = \{0, 1, \ldots, |q| - 1\}$ and the difference set $\Delta D := D - D$, then the CC digit set $\mathcal{D} = Dv$ and $\Delta\mathcal{D} := \mathcal{D} - \mathcal{D} = \Delta Dv$. Without loss of generality, we can assume the digit set $\mathcal{D}$ is primitive [LW2] (i.e., $\mathbb{Z}^2 = \{\gamma v + \delta Av : \gamma, \delta \in \mathbb{Z}\}$) and $T + \mathbb{Z}^2$ is a lattice tiling. It is easy to see that $T + \ell$ is a neighbor of T if and only if

$$\ell = \gamma v + \delta Av = \sum_{i=1}^{\infty} b_i A^{-i} v \in T - T, \text{ where } b_i \in \Delta D.$$

The following is a neighbor-generating formula which plays a key role in constructing the labeled directed graph for the boundary.

Lemma 2.1. *(*[LL]*) Suppose $T + \ell$ is a neighbor of T with $\ell = \gamma v + \delta Av = \sum_{i=1}^{\infty} b_i A^{-i} v$, then we get another neighbor $T + \ell'$ satisfying $\ell' = A\ell - b_1 v = \gamma' v + \delta' Av$ with $\gamma' = -(q\delta + b_1)$ and $\delta' = \gamma - p\delta$.*

Inductively, we can construct a sequence of neighbors: $\{T + \ell_n\}_{n=0}^{\infty}$ where $\ell_0 = \ell$ and $\ell_{n+1} = A\ell_n - b_{n+1}v$.

Let T be a disk-like CC tile and $\mathcal{V} = \{\ell : T + \ell \text{ is a neighbor of } T\}$. Let $T_\ell = T \cap (T + \ell)$. Then the boundary of T can be written as

$$\partial T = \bigcup_{\ell \in \mathcal{V}} T_\ell. \tag{2.1}$$

Define an edge set $\mathcal{E} := \{e = (\ell, \ell') : \ \ell, \ \ell' \in \mathcal{V} \ \text{ and } \ \ell' = A\ell - b_1 v \text{ for some } b_1 \in \Delta D\}$ and a labeling $\mathcal{L} : \mathcal{E} \to \mathcal{A}$ by $\mathcal{L}(e) = b_1$ where $\mathcal{A} = \Delta D$. Then by the definition above, $\mathbf{G} = (\mathcal{G}, \mathcal{L})$ is a labeled directed graph and it determines a sofic shift. We call $\mathbf{G}$ the *neighbor graph* of T.

Proposition 2.2. *Let* $\mathbf{G}$ *be the neighbor graph of a CC disk-like tile* T. *If* $x = \sum_{i=1}^{\infty} a_i A^{-i} v = \ell + \sum_{i=1}^{\infty} a'_i A^{-i} v \in T_\ell$ *where* $a_i, a'_i \in D$, *then* $\{b_i := a_i - a'_i\}_{i=1}^{\infty}$ *is the sequence of labeling of the edges of an infinite path starting at* ℓ *(or simply called a label sequence starting at* ℓ*). Conversely, any label sequence* $\{b_i\}_{i=1}^{\infty}$ *(with* $b_i \in \Delta D$*) starting at* ℓ *defines a set*

$$\Big\{x : x = \sum_{i=1}^{\infty} a_i A^{-i} v = \ell + \sum_{i=1}^{\infty} a'_i A^{-i} v,\ a_i - a'_i = b_i,\ a_i, a'_i \in D \text{ for } i = 1, 2, \ldots\Big\}$$

of boundary points of T.

Proof. Since $\ell = \sum_{i=1}^{\infty} b_i A^{-i} v$ with $b_i = a_i - a'_i$, by Lemma 2.1, we have a sequence of neighbors $\{T + \ell_n\}_{n=0}^{\infty}$ where $\ell_0 = \ell$ and $\ell_{n+1} = A\ell_n - b_{n+1} v$, hence $\{b_i\}_{i=1}^{\infty}$ is a label sequence starting at ℓ by the definition.

Conversely, if $\ell = \sum_{i=1}^{\infty} b_i A^{-i} v$ where $b_i \in \Delta D$, then $b_i = a_i - a'_i$ for $a_i, a'_i \in D$ and $\ell = \sum_{i=1}^{\infty} (a_i - a'_i) A^{-i} v$. It follows that

$$x = \sum_{i=1}^{\infty} a_i A^{-i} v = \ell + \sum_{i=1}^{\infty} a'_i A^{-i} v \in T \cap (T + \ell) = T_\ell. \tag{2.2}$$

□

We can verify whether the origin 0 is a boundary point of T by the following method.

Corollary 2.3. $0 \in \partial T$ *if and only if there exists an infinite path in* $\mathbf{G}$ *with all edge labels either non-positive or non-negative.*

Proof. Suppose $0 \in T \cap (T + \ell)$ for some neighbor $T + \ell$. Putting $a_i = 0$ for all i into (2.2), we have

$$\ell = \sum_{i=1}^{\infty} (-a'_i) A^{-i} v.$$

Since $a'_i \in D$, the label sequence $\{b_i = -a'_i\}_{i=1}^{\infty}$ starting at ℓ has all labels non-positive. Similarly $\{b'_i = a_i\}_{i=1}^{\infty}$ is a sequence starting at $-\ell$ with all labels non-negative. By reversing the argument, we can prove the converse. □

In fact, we can determine the neighbor graph $\mathbf{G}$ for any disk-like CC tile T. Let us take the case of $f(x) = x^2 + px + q,\ p, q \geq 2$ (excluding $p = q = 2$) as an example. By Theorem 1.1, T is a hexagonal tile with six neighbors [LL] and

$$\mathcal{V} = \{\pm v,\ \pm\big(Av + (p-1)v\big),\ \pm(Av + pv)\}. \tag{2.3}$$

In view of the definition of $\mathcal{E}$, if $\ell = v$ we take $b_1 = -p$ and $\ell' = Av + pv$ or $b_1 = -(p-1)$ and $\ell' = Av + (p-1)v$; if $\ell = Av + pv$, using $f(A)v = 0$, we have $b_1 = -(q-1)$ and $\ell' = -v$. Proceeding similarly with all ℓ, we obtain Table 1. Then

ℓ	b_1	ℓ'
v	$-(p-1)$	$Av+(p-1)v$
	$-p$	$Av+pv$
$Av+(p-1)v$	$-(q-p)$	$-Av-pv$
	$-(q-p+1)$	$-Av-(p-1)v$
$Av+pv$	$-(q-1)$	$-v$
$-v$	$p-1$	$-Av-(p+1)v$
	p	$-Av-pv$
$-Av-(p-1)v$	$q-p$	$Av+pv$
	$q-p+1$	$Av+(p-1)v$
$-Av-pv$	$q-1$	v

TABLE 1. Relation among all neighbors of T associated with $f(x) = x^2+px+q,\ p,q \geq 2,\ 2p \leq q+2$ (excluding $p=q=2$).



FIGURE 1. The neighbor graph of T associated with $f(x) = x^2+px+q,\ p,q \geq 2,\ 2p \leq q+2$, (excluding $q=p=2$).

we establish the neighbor graph (Figure 1). The neighbor graphs corresponding to other $f(x)$ are given in Appendix A.

Following [LW1], we let $\mathcal{D}_{A,k} = \{\sum_{i=0}^{k-1} a_i A^i v : a_i \in D\}$, $\Delta\mathcal{D}_{A,k} = \mathcal{D}_{A,k} - \mathcal{D}_{A,k} = \{\sum_{i=0}^{k-1} b_i A^i v : b_i \in \Delta D\}$ and $\mathcal{D}_{A,\infty} = \cup_{k=1}^{\infty}\mathcal{D}_{A,k}$.

Proposition 2.4. *Let T be a disk-like CC tile and $T+\ell$ a neighbor. Then $\ell = \sum_{i=0}^{k} b_i A^i v \in \Delta\mathcal{D}_{A,k+1}$ for some $k \in \mathbb{Z}$ with $b_k \in \{-1,1\}$, $b_i \in \Delta D$ for $0 \leq i < k$. When $f(x) = x^2 \pm 2x + 2$, $k=3$; and $k=1$ otherwise.*

Proof. It follows from (2.3) that $\ell \in \Delta\mathcal{D}_{A,2}$ excluding the case of $f(x) = x^2 \pm 2x + 2$. For $f(x) = x^2 \pm 2x + 2$, we have $Av \pm 2v = A^3v \pm A^2v + Av \in \Delta\mathcal{D}_{A,4}$ by using $(A \mp I)f(A) = 0$. $\square$

A more desirable property is that any $\ell \in \mathbb{Z}^2$ can be expressed as $\ell = \sum_{i=0}^{k} a_i A^i v \in \mathcal{D}_{A,k+1}$ (instead of $\Delta\mathcal{D}_{A,k+1}$) for some $k \in \mathbb{Z}$ with $a_k = 1$ and $a_i \in D$ where $0 \leq i < k$. But this is not always the case. This property is closely related to a *number system* defined below (see also [MTT]).

Definition 2.5. *Let $A \in M_2(\mathbb{Z})$ be expanding and $\mathcal{D}$ be a CC digit set. The self-affine pair $(A, \mathcal{D})$ is said to be a number system if for any $\ell \in \mathbb{Z}^2$, it has a unique representation $\ell = \sum_{i=0}^{k} A^i v_i'$ with $v_i' \in \mathcal{D}$.*

For convenience, we sometimes write a point of the form $x = \sum_{i=1}^{\infty} a_i A^{-i} v \in \mathbb{R}^2$ as *radix expansion*: $0.a_1a_2a_3\ldots$ An overbar denotes repeating digits as in $0.12\overline{301} = 0.12301301301\ldots$ Likewise, $a_{-2}a_{-1}a_0.a_1a_2a_3\ldots$ represents a point $a_{-2}A^2v + a_{-1}Av + a_0v + \sum_{i=1}^{\infty} a_i A^{-i} v$. Note that shifting a radix place to the left means multiplying A to x. When x is on the boundary of T, the radix expansion of x is not unique. Now we give some equivalent conditions for the self-affine pair $(A, \mathcal{D})$ to be a number system.

Theorem 2.6. *Let $T = T(A, \mathcal{D})$ be a disk-like CC tile. Then the following are equivalent:*

(i) $(A, \mathcal{D})$ is a number system.

(ii) $0 \in T^\circ$.

(iii) $f(x) = x^2 + px + q$ with $-1 \leq p$ and $q \geq 2$.

(iv) For all neighbors $T + \ell$, $\ell = \sum_{i=0}^{k} a_i A^i v \in \mathcal{D}_{A,k+1}$ for some $k \in \mathbb{Z}$ with $a_k = 1$ and $a_i \in D$ where $0 \leq i < k$.

Proof. (i)⇒(ii) Suppose $0 \notin T^\circ$. Then $0 \in T \cap (T + \ell)$ for some $\ell \in \mathbb{Z}^2 \setminus \{0\}$. Since $(A, \mathcal{D})$ is a number system, $\ell = \sum_{i=-k}^{0} a_i A^{-i} v$ with $a_i \in D$ and $a_{-k} > 0$. Hence $0 = a_{-k}a_{-(k-1)}\ldots a_{-1}a_0.a_1a_2a_3\ldots$. Shifting the radix point k places to the left, we get $0 = a_{-k}.a_{-(k-1)}\ldots a_{-1}a_0a_1a_2a_3\ldots$. That means $T + a_{-k}v$ is a neighbor of T. By Proposition 2.4, $a_{-k} = 1$. Hence 0 corresponds to an infinite path starting at v with non-positive labels $b_i = -a_i$. But by checking all the neighbor graphs in Appendix A, we find no such path.

(ii)⇒(i) It suffices to show that $\mathbb{Z}^2 \subset \mathcal{D}_{A,\infty}$. By the lattice tiling property, 0 is the only lattice point in T, i.e., $\mathbb{Z}^2 \cap T = \{0\}$. It follows that $\mathbb{Z}^2 \cap A^nT = \sum_{i=0}^{n-1} A^i\mathcal{D} = \mathcal{D}_{A,n}$ for $n \geq 1$. If $\ell \in \mathbb{Z}^2$, there exists a large integer n such that $\ell \in A^nT$ as $0 \in T^\circ$, then $\ell \in \mathcal{D}_{A,n} \subset \mathcal{D}_{A,\infty}$.

(ii)⇔(iii) By inspecting all neighbor graphs in Appendix A, we find that in each graph corresponding to $f(x) = x^2 + px + q$ with $-1 \leq p$ and $q \geq 2$, there exists no infinite path with edge labels either all non-positive or all non-negative, hence $0 \in T^\circ$ by Corollary 2.3. In every other case, there always exists such a path. All these paths are listed in Table 2.

$f(x)$	Neighbor	Path
x^2-2x+2	$Av-v$	$\overline{(-1)}$
	$-Av+v$	$\overline{1}$
x^2-px-q	v	$\overline{p(q-1)}$
	$-v$	$\overline{(-p)[-(q-1)]}$
x^2+px-q	$Av+(p+1)v$	$\overline{(q-p-1)}$
	$-Av-(p+1)v$	$\overline{[-(q-p-1)]}$
x^2-q	v	$\overline{0(q-1)}$
	$-v$	$\overline{0[-(q-1)]}$
	$Av+v$	$\overline{(q-1)}$
	$-Av-v$	$\overline{[-(q-1)]}$
x^2-px+q	$Av-(p-1)v$	$\overline{[-(q-p+1)]}$
	$-Av+(p-1)v$	$\overline{(q-p+1)}$

TABLE 2. Infinite paths representing a boundary point 0.

(iii)⇒(iv) Let $f(x)$ be one of the cases: x^2+q, x^2+x+q, x^2+px+q ($p \geq 2$, excluding $p=q=2$), x^2+2x+2, x^2-x+q, where $p \geq 0$ and $q \geq 2$. In each case, we can rewrite their neighbors as the desired form in (iv). By using $0=f(A)v$, $0=(A-I)f(A)v$, $0=(A+I)f(A)v$, we have

Case (1) $f(x)=x^2+q$. $Av-v=A^2v+Av+(q-1)v$, $-v=A^2v+(q-1)v$, $-Av=A^3v+(q-1)Av$, $-Av+v=A^3v+(q-1)Av+v$, $-Av-v=A^3v+A^2v+(q-1)Av+(q-1)v$.

Case (2) $f(x)=x^2+x+q$. $-v=A^2v+Av+(q-1)v$, $-Av=A^3v+A^2v+(q-1)Av$, $-Av-v=A^2v+(q-1)v$.

Case (3) $f(x)=x^2+px+q$ ($p \geq 2$). $-v=A^2v+pAv+(q-1)v$, $-Av-(p-1)v=A^2v+(p-1)Av+(q-p+1)v$, $-Av-pv=A^2v+(p-1)Av+(q-p)v$.

Case (4) $f(x)=x^2+2x+2$. $Av+2v=A^3v+A^2v+Av$, $-v=A^4v+A^3v+A^2v+v$, $-Av-v=A^2v+Av+v$, $-Av-2v=A^2v+Av$.

Case (5) $f(x)=x^2-x+q$. $Av-v=A^2v+(q-1)v$, $-v=A^3v+(q-1)Av+(q-1)v$, $-Av=A^4v+(q-1)A^2v+(q-1)Av$, $-Av+v=A^4v+(q-1)A^2v+(q-1)Av+v$.

(iv)⇒(ii) Suppose $0 \notin T^{\circ}$. By the same argument as in the proof of (i)⇒(ii) above, there should be an infinite path in the neighbor graph starting at v with edge labels all non-positive. But we find no such paths by inspecting all the neighbor graphs in Appendix A. $\square$

Remark 2.7. *Gilbert* [Gi] *obtained some related results in the context of quadratic number fields. We conjecture that Theorem 2.6 can be extended to non-disk-like tiles.*

3. **Boundary of T as Graph-directed Sets**

For a directed graph $\mathcal{G} = \mathcal{G}(\mathcal{V}, \mathcal{E})$ where $\mathcal{V} = \{v_1, v_2, \ldots, v_m\}$, we write $\mathcal{E}_{i,j}$ for the set of edges from vertex v_i to vertex v_j, and we add a contraction mapping $F_e : \mathbb{R}^2 \to \mathbb{R}^2$ for each edge $e \in \mathcal{E}$. Then the family of contractions $\{F_e : e \in \mathcal{E}\}$ is called a *graph-directed iterated function system (graph-directed IFS)* and it is well-known that there exists a unique family of non-empty compact subsets $E_1, \ldots, E_m$ of $\mathbb{R}^2$ ([Fa2], [MW]) such that

$$E_i = \bigcup_{j=1}^{m} \bigcup_{e \in \mathcal{E}_{i,j}} F_e(E_j). \tag{3.1}$$

We call $\{E_1, \ldots, E_m\}$ a family of *graph-directed sets.* The graph-directed IFS $\{F_e : e \in \mathcal{E}\}$ is said to satisfy the *open set condition (OSC)* if there exist a family of open sets $\{O_1, \ldots, O_m\}$ such that

$$O_i \supset \bigcup_{j=1}^{m} \bigcup_{e \in \mathcal{E}_{i,j}} F_e(O_j) \quad \text{for } i = 1, 2, \ldots, m \tag{3.2}$$

with disjoint unions, i.e., $F_e(O_j) \cap F_{e'}(O_{j'}) = \emptyset$ whenever $(e, j) \neq (e', j')$.

In this section, we will show that the relationship of $\{T_\ell\}_{\ell \in \mathcal{V}}$ where $\mathcal{V} = \{\ell : T + \ell \text{ is a neighbor of } T\}$ as in Section 2 can be expressed as a family of graph-directed sets by means of the following proposition, which applies to non-disk-like T as well.

Proposition 3.1. *Let $\ell = \gamma v + \delta A v$, $\ell' = \gamma' v + \delta' A v \in \mathcal{V}$ such that $\ell' = A\ell - b_1 v$ for some $b_1 \in \Delta D$, then*

$$A^{-1}(T_{\ell'} + jv) \subset T_\ell \quad \textit{for all} \quad j \in I_{b_1} := \begin{cases} \{b_1, b_1 + 1, \ldots, q-1\} & \textit{if} \quad b_1 \geq 0; \\ \{0, 1, \ldots, q - 1 + b_1\} & \textit{if} \quad b_1 < 0. \end{cases}$$

Moreover,

$$T_\ell = \bigcup_{\ell' \in B_\ell} \bigcup_{j \in I_{b_1}} A^{-1}(T_{\ell'} + jv)$$

where $B_\ell := \{\ell'' \in \mathcal{V} : \ell'' = A\ell - b_1' v \text{ for some } b_1' \in \Delta D\}$.

Proof. When $b_1 \geq 0$, if $x \in T_{\ell'}$ then the radix expansion is

$$x = 0.c_1c_2c_3\ldots = \delta'\gamma'.c_1'c_2'c_3'\ldots.$$

It follows from Lemma 2.1 and $0 = A^{-1}f(A)v$ that

$$A^{-1}x + (b_1 + k)A^{-1}v = 0.(b_1 + k)c_1c_2c_3\ldots = \delta\gamma.kc_1'c_2'c_3'\ldots \in T_\ell$$

for $k = 0, 1, \ldots, q - 1 - b_1$. The case when $b_1 < 0$ can be proved similarly.

For the second part, we only need to show

$$T_\ell \subset \bigcup_{\ell' \in B_\ell} \bigcup_{j \in I_{b_1}} A^{-1}(T_{\ell'} + jv).$$

Let $y = 0.a_1a_2a_3\ldots = \delta\gamma.a_1'a_2'a_3'\ldots \in T_\ell$. It follows that $Ay - a_1v = 0.a_2a_3a_4\ldots = \delta\gamma(a_1' - a_1).a_2'a_3'\ldots \in T_{\ell'}$, where $\ell' = A\ell - (a_1 - a_1')v$. This implies $y \in A^{-1}(T_{\ell'} + a_1v)$. By definition, we see that $a_1 \in I_{b_1}$ for $b_1 = a_1 - a_1'$. $\square$

Therefore, ∂T is the union of a family of graph-directed sets. We give the case $f(x) = x^2 + px + q$ ($p, q \geq 2$ excluding $p = q = 2$) as an example in the following. All the other cases are given in Appendix B.

Example 3.2. *Consider the case $f(x) = x^2 + px + q$ ($p, q \geq 2,$ excluding $p = q = 2$). When $\ell = v$, from Table 1 we have $B_\ell = B_v = \{Av + pv, Av + (p-1)v\}$. When $\ell' = Av + pv$, $b_1 = -p$ and $I_{-p} = \{0, 1, 2, \ldots, q-1-p\}$; when $\ell' = Av + (p-1)v$, $b_1 = -(p-1)$ and $I_{-(p-1)} = \{0, 1, 2, \ldots, q-p\}$. Thus by Proposition 3.1, the first set equation comes out. Similarly the other five can be deduced. For simplicity we let $u_1 = v$, $u_2 = Av + (p-1)v$, $u_3 = Av + pv$. Then the graph-directed sets $T_{\pm u_1}$, $T_{\pm u_2}$, $T_{\pm u_3}$, representing ∂T satisfy*

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=0}^{q-p}(T_{u_2} + jv) \cup \bigcup_{j=0}^{q-p-1}(T_{u_3} + jv) \\
AT_{u_2} &= \bigcup_{j=0}^{p-2}(T_{-u_2} + jv) \cup \bigcup_{j=0}^{p-1}(T_{-u_3} + jv) \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= \bigcup_{j=p-1}^{q-1}(T_{-u_2} + jv) \cup \bigcup_{j=p}^{q-1}(T_{-u_3} + jv) \\
AT_{-u_2} &= \bigcup_{j=q-p+1}^{q-1}(T_{u_2} + jv) \cup \bigcup_{j=q-p}^{q-1}(T_{u_3} + jv) \\
AT_{-u_3} &= T_{u_1} + (q-1)v
\end{aligned}$$

Theorem 3.3. *The graph-directed IFS representing the boundary of a disk-like CC tile T satisfies the OSC.*

Proof. Replacing T_ℓ by $(T + \ell)^\circ$, we can check that the graph-directed IFS representing ∂T satisfies the OSC. We illustrate the idea by proving the case $f(x) = x^2 + px + q$ ($p \geq 2, q \geq 2,$ excluding $p = q = 2$). In view of Example 3.2, we need to show

$$\begin{aligned}
A(T+u_1)^\circ &\supset \bigcup_{j=0}^{q-p}\big((T+u_2)^\circ+jv\big)\cup\bigcup_{j=0}^{q-p-1}\big((T+u_3)^\circ+jv\big)\\
A(T+u_2)^\circ &\supset \bigcup_{j=0}^{p-2}\big((T-u_2)^\circ+jv\big)\cup\bigcup_{j=0}^{p-1}\big((T-u_3)^\circ+jv\big)\\
A(T+u_3)^\circ &\supset (T-u_1)^\circ\\
A(T-u_1)^\circ &\supset \bigcup_{j=p-1}^{q-1}\big((T-u_2)^\circ+jv\big)\cup\bigcup_{j=p}^{q-1}\big((T-u_3)^\circ+jv\big)\\
A\big(T-u_2\big)^\circ &\supset \bigcup_{j=q-p+1}^{q-1}\big((T+u_2)^\circ+jv\big)\cup\bigcup_{j=q-p}^{q-1}\big((T+u_3)^\circ+jv\big)\\
A(T-u_3)^\circ &\supset (T+u_1)^\circ+(q-1)v
\end{aligned}$$

with disjoint unions. Since T is a CC tile, it follows that

$$AT^\circ\supset\bigcup_{j=0}^{q-1}(T+jv)^\circ=\bigcup_{j=0}^{q-1}(T^\circ+jv) \tag{3.3}$$

with disjoint union. By using (3.3) and $0=f(A)v=A^2v+pAv+qv$ extensively, we prove the first two set inequalities in the following. The remaining four can be verified similarly.

For $j=0,1,\dots,q-p$,

$$\big(T+u_2\big)^\circ+jv=T^\circ+(p-1+j)v+Av\subset A(T+u_1)^\circ.$$

For $j=0,1,\dots,q-p-1$,

$$(T+u_3)^\circ+jv=T^\circ+(p+j)v+Av\subset A(T+u_1)^\circ.$$

For $j=0,1,\dots,p-2$,

$$\begin{aligned}
\big(T-u_2\big)^\circ+jv &= T^\circ+(j-p+1)v-Av\\
&= T^\circ+(q+j-p+1)v+A^2v+(p-1)Av\\
&\subset A(T+u_2)^\circ.
\end{aligned}$$

For $j=0,1,\dots,p-1$,

$$\begin{aligned}
(T-u_3)^\circ+jv &= T^\circ+(j-p)v-Av\\
&= T^\circ+(q+j-p)v+A^2v+(p-1)Av\\
&\subset A(T+u_2)^\circ.
\end{aligned}$$

By the same way, all the other cases follow and hence the theorem is proved. □

The Hausdorff dimension ($\dim_H$) (see e.g., [Fa], [Fa2]) is the most common and important dimension in fractal geometry. Since a disk-like tile has positive Lebesgue measure, its Hausdorff dimension is simply the dimension of the underlying space

$\mathbb{R}^2$. A more interesting problem is to find the Hausdorff dimension of the boundary of the T. In the rest of this section, we find the exact value of $\dim_H(\partial T)$ for certain particular cases that A is a similarity. We state the simplest one first.

Proposition 3.4. *Let* $A \in M_2(\mathbb{Z})$ *be expanding with characteristic polynomial* $f(x) = x^2 + q$ $(|q| \geq 2)$ *and* $T(A, \mathcal{D})$ *a disk-like CC tile. Then* $\dim_H(\partial T) = 1$.

Proof. By Theorem 1.1, T is a square tile (parallelogram). Hence $\dim_H(\partial T) = 1$. ☐

Geometrically, a similarity is a multiple of either a reflection or a rotation. We call the former a *scaled reflection* and the latter a *scaled rotation*; algebraically, a similarity is a multiple of an orthogonal matrix. The case that A is a scaled reflection is solved already as its characteristic polynomial is of the form $f(x) = x^2 - q$ $(q > 0)$. So we focus our attention on those A that are scaled rotations.

Lemma 3.5. *Let* A *be a scaled rotation. Then its characteristic polynomial has positive constant term and* A *has either two distinct non-real eigenvalues or two equal real eigenvalues.*

Proof. Let $A = \begin{pmatrix} r\cos\theta & -r\sin\theta \\ r\sin\theta & r\cos\theta \end{pmatrix}$. The characteristic polynomial is given by $x^2 - 2r\cos\theta\, x + r^2$. It has two equal real zeros when $\theta = 0$ or π and two distinct non-real zeros otherwise. ☐

We shall make use of the following well-known dimension formula (see also [DKV], [Fa2], [HLR], [LN], [MW], [SW]) to compute the Hausdorff dimension of ∂T.

Theorem 3.6. *If* A *is a similarity with* $|\det(A)| = |q| \geq 2$, *then the Hausdorff dimension of* ∂T *is given by*

$$\dim_H(\partial T) = \log \rho(M)/\log r = 2\log \rho(M)/\log |q|, \tag{3.4}$$

where $\rho(M)$ *denotes the spectral radius of the contact matrix* M *and* $r = |q|^{1/2}$ *is the expansion ratio of* A.

Remark 3.7. *In* [MW], *a separation condition is required for the validity of (3.4). It is relaxed to the OSC in* [Fa2]. *It is shown in* [DKV], [HLR], [LN] *and* [SW] *that (3.4) is also valid with a weaker assumption of the OSC.*

We first find the *contact matrix (or substitution matrix)* M for T. Let ℓ, ℓ', b_1 and B_ℓ be defined as in Proposition 3.1. The entry $M_{\ell\ell'}$ is defined by $M_{\ell\ell'} := \#\big((A\ell + \mathcal{D}) \cap (\ell' + \mathcal{D})\big)$. Since $\mathcal{D}$ is a CC digit set, it can be shown easily that $M_{\ell\ell'} = \#I_{b_1} = q - |b_1|$ where I_{b_1} is as in Proposition 3.1. Recall that b_1 is the label of the edge starting at ℓ and ending at ℓ'. Hence we obtain the contact matrix M of T from its neighbor graph with different edge labels (i.e., replace b_1 by $q - |b_1|$). There is a one-to-one correspondence between the contact matrix and the neighbor graph. For example, the contact matrix for the case $f(x) = x^2 + px + q$ $(p, q \geq 2,\ 2p \leq q + 2$ excluding $p = q = 2)$ can be found in Table 3, and the related

	v	$Av+(p-1)v$	$Av+pv$	$-v$	$-Av-(p-1)v$	$-Av-pv$
v	0	$q-p+1$	$q-p$	0	0	0
$Av+(p-1)v$	0	0	0	0	$p-1$	p
$Av+pv$	0	0	0	1	0	0
$-v$	0	0	0	0	$q-p+1$	$q-p$
$-Av-(p-1)v$	0	$p-1$	p	0	0	0
$-Av-pv$	1	0	0	0	0	0

TABLE 3. The contact matrix of T associated with $f(x) = x^2 + px + q,\ p, q \geq 2, 2p \leq q + 2$ (excluding $p = q = 2$).

neighbor graph is shown by Figure 1. The contact matrices for the other cases are given in Appendix C.

If M is irreducible (i.e., for each entry M_{ij}, there exists an integer $n \geq 0$ such that $(M^n)_{ij} > 0$), then the spectral radius $\rho(M) = \lambda_M$ where λ_M is the *Perron-Frobenius eigenvalue* of M as stated in the following simplified version of the Perron-Frobenius Theorem (see [Ga] or [Se] for details).

Theorem 3.8. *Let M be an irreducible non-negative matrix. Then there exists a positive eigenvalue λ_M such that $\lambda_M \geq |\mu|$ for all eigenvalues μ of M. Moreover, λ_M is a simple zero of the characteristic polynomial of M.*

It is known that a contact matrix is irreducible if and only if the neighbor graph it represents is strongly connected. A directed graph is called *strongly connected* if for any two vertices v_i, v_j there exists a path starting at v_i and ending at v_j. Now we state and prove our main result of this section.

Theorem 3.9. *Let $A \in M_2(\mathbb{Z})$ be an expanding similarity with characteristic polynomial $f(x) = x^2 + px + q$ and $T(A, \mathcal{D})$ be a disk-like CC tile. Then $\rho(M)$ is the largest positive zero of the cubic polynomial*

$$x^3 - (|p| - 1)x^2 - (|q| - |p|)x - |q|.$$

Hence $\dim_H \big(\partial T\big) = 2 \log \rho(M) / \log |q|$.

Proof. Since $|\det(A)| = |q|$, it is more convenient to work with $f(x) = x^2 \pm px \pm q$ $(p \geq 0,\ q \geq 2)$. Also we ignore those $f(x)$ of the form $f(x) = x^2 \pm px - q$ $(p > 0,\ q \geq 2)$ as they cannot be characteristic polynomials of similarities (Lemma 3.5). In view of Appendix A or C, we can check that the contact matrix is irreducible if $f(x)$ is one of $x^2 \pm 2x + 2,\ x^2 \pm x + q,\ x^2 \pm px + q$ $(p \geq 2$ excluding $p = q = 2)$; and is reducible if $f(x) = x^2 \pm q$. Hence we can apply Theorem 3.8 for all the cases except $f(x) = x^2 \pm q$.

Case (1) $f(x) = x^2 + 2x + 2$. The characteristic polynomial of the corresponding contact matrix M is $(x - 1)(x^2 + 2x + 2)(x^3 - x^2 - 2)$. We see that $\rho(M)$ is equal to the positive zero of $x^3 - x^2 - 2$. By the cubic equation formula, we have

$$\rho(M) = \frac{[28 + (28^2 - 1)^{1/2}]^{1/3} + [28 - (28^2 - 1)^{1/2}]^{1/3}}{3} \approx 1.69562.$$

Case (2) $f(x) = x^2 - 2x + 2$. The characteristic polynomial of the corresponding M is $(x+1)(x^2 - 2x + 2)(x^3 - x^2 - 2)$. We have the same $\rho(M)$ as in the case $f(x) = x^2 + 2x + 2$.

Case (3) $f(x) = x^2 + x + q$. The characteristic polynomial of the corresponding M is $(x-1)(x^2 + x + q)[x^3 - (q-1)x - q]$. Notice that $\rho(M) \neq 1$. If $\rho(M) = 1$, then $\dim_{\mathcal{H}}(\partial T) = 0$, which implies ∂T is totally disconnected (Proposition 2.5, [Fa]). This is not possible for the boundary of a topological disk. The zeros of $x^2 + x + q$ are non-real. So $\rho(M)$ is the largest positive real zero of $x^3 - (q-1)x - q$.

Case (4) $f(x) = x^2 - x + q$. The characteristic polynomial of the corresponding M is $(x+1)(x^2 - x + q)[x^3 - (q-1)x - q]$. The zeros of $x^2 - x + q$ are non-real. So, as in the previous case, $\rho(M)$ is the largest positive real zero of $x^3 - (q-1)x - q$.

Case (5) $f(x) = x^2 + px + q$. The characteristic polynomial of the corresponding M is $(x-1)(x^2 + px + q)[x^3 - (p-1)x^2 - (q-p)x - q]$. As explained before, $\rho(M) \neq 1$. The zeros of $x^2 + px + q$ are either both negative or both non-real. Hence $\rho(M)$ is the largest positive real zero of $x^3 - (p-1)x^2 - (q-p)x - q$.

Case (6) $f(x) = x^2 - px + q$. The characteristic polynomial of the corresponding M is $(x+1)(x^2 - px + q)[x^3 - (p-1)x^2 - (q-p)x - q]$. We know that $f(x)$ cannot have unequal real zeros. So we must have $p^2 - 4q \leq 0$. When $p^2 - 4q < 0$, the zeros of $x^2 - px + q$ are non-real. Then $\rho(M)$ is the largest positive real zero of $x^3 - (p-1)x^2 - (q-p)x - q$. When $p^2 - 4q = 0$, the two zeros of $x^2 - px + q$ are equal. But the Perron-Frobenius eigenvalue should be a simple zero of the characteristic polynomial of M (Theorem 3.8), so $\rho(M)$ is also the largest positive real zero of $x^3 - (p-1)x^2 - (q-p)x - q$.

Case (7) $f(x) = x^2 + q$. The contact matrix M is reducible. Its characteristic polynomial is $(x^2 - q)(x^2 + q)(x-1)(x+1)(x^2+1)$. We see that $\rho(M) = q^{1/2}$, which is also the largest positive zero of $x^3 + x^2 - qx - q = (x^2 - q)(x+1)$.

Case (8) $f(x) = x^2 - q$. The contact matrix M is also reducible and its characteristic polynomial is found to be $(x^2 - q)^2(x+1)(x-1)^3$. As in the previous case, $\rho(M) = q^{1/2}$, which is also the largest positive zero of $x^3 + x^2 - qx - q$. $\square$

Remark 3.10. *It is interesting to see that the signs of* p *and* q *do not matter in the calculation of* $\dim_H(\partial T)$ *when* A *is a similarity. Notice also for the last two cases,* $f(x) = x^2 + q$ $(|q| \geq 2)$, *we have* $\rho(M) = |q|^{1/2}$. *It follows that* $\dim_H(\partial T) = 1$, *as expected for the boundary of a parallelogram (Proposition 3.4).*

We observe that $\dim_H(\partial T)$ is independent of the choice of the vector v in the following sense.

Corollary 3.11. *Let* $A \in M_2(\mathbb{Z})$ *be an expanding similarity with characteristic polynomial* $f(x) = x^2 + px + q$ $(|q| \geq 2)$. *Let* $\mathcal{D} = \mathcal{D}(v, |q|)$ *and* $\mathcal{D}' = \mathcal{D}(v', |q|)$ *be two CC digit sets such that each of* $\{v, Av\}$ *and* $\{v', Av'\}$ *is an independent set. If* $2|p| \leq |q+2|$, *then*

$$\dim_H\big(\partial T(A, \mathcal{D})\big) = \dim_H\big(\partial T(A, \mathcal{D}')\big).$$

Proof. As $2|p| \leq |q+2|$, both $T(A, \mathcal{D})$ and $T(A, \mathcal{D}')$ are disk-like CC tiles (Theorem 1.1). Hence the corollary follows from Theorem 3.9. $\square$

Remark 3.12. *We conjecture that Theorem 3.9 and Corollary 3.11 are also valid when* $2|p| > |q+2|$, *i.e.,* T *is non-disk-like. The major difficulty in justifying these conjectures is that, in general, there is no upper bound on the number of neighbors of a non-disk-like CC tile* [DJN].

4. Appendix A: Neighbor Graphs

Let $f(x) = x^2 \pm px \pm q$ $(p \geq 0,\ q \geq 2)$. The neighbor graphs of disk-like tiles are classified by $f(x)$ and listed below.



(a) $f(x) = x^2 + q$ (b) $f(x) = x^2 - q$

Figure 2.



(a) $f(x) = x^2 + x + q$ (b) $f(x) = x^2 - x + q$

Figure 3.



(a) $f(x) = x^2 + px + q$, $p \geq 2$, $2p \leq q + 2$ (excluding $q = p = 2$)

(b) $f(x) = x^2 - px + q$, $p \geq 2$, $2p \leq q + 2$ (excluding $q = p = 2$)

FIGURE 4.



(a) $f(x) = x^2 + px - q$, $p \geq 1$, $2p \leq q - 2$

(b) $f(x) = x^2 - px - q$, $p \geq 1$, $2p \leq q - 2$

FIGURE 5.



(a) $f(x) = x^2 + 2x + 2$

(b) $f(x) = x^2 - 2x + 2$

FIGURE 6.

5. **Appendix B: Graph-directed Sets**

Let $f(x) = x^2 \pm px \pm q$ $(p \geq 0,\ q \geq 2)$. The graph-directed sets representing the boundary ∂T are classified by $f(x)$ and listed below.

(1) $f(x) = x^2 + q$. Convention: $u_1 = v$, $u_2 = Av - v$, $u_3 = Av$, $u_4 = Av + v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=1}^{q-1}(T_{u_2}+jv) \cup \bigcup_{j=0}^{q-1}(T_{u_3}+jv) \cup \bigcup_{j=0}^{q-2}(T_{u_4}+jv)\\
AT_{u_2} &= T_{-u_4}\\
AT_{u_3} &= T_{-u_1}\\
AT_{u_4} &= T_{u_2}\\
AT_{-u_1} &= \bigcup_{j=0}^{q-2}(T_{-u_2}+jv) \cup \bigcup_{j=0}^{q-1}(T_{-u_3}+jv) \cup \bigcup_{j=1}^{q-1}(T_{-u_4}+jv)\\
AT_{-u_2} &= T_{u_4}+(q-1)v\\
AT_{-u_3} &= T_{u_1}+(q-1)v\\
AT_{-u_4} &= T_{-u_2}+(q-1)v
\end{aligned}$$

(2) $f(x) = x^2 - q$. Convention: $u_1 = v$, $u_2 = Av - v$, $u_3 = Av$, $u_4 = Av + v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=1}^{q-1}(T_{u_2}+jv) \cup \bigcup_{j=0}^{q-1}(T_{u_3}+jv) \cup \bigcup_{j=0}^{q-2}(T_{u_4}+jv)\\
AT_{u_2} &= T_{-u_2}+(q-1)v\\
AT_{u_3} &= T_{u_1}+(q-1)v\\
AT_{u_4} &= T_{u_4}+(q-1)v\\
AT_{-u_1} &= \bigcup_{j=0}^{q-2}(T_{-u_2}+jv) \cup \bigcup_{j=0}^{q-1}(T_{-u_3}+jv) \cup \bigcup_{j=1}^{q-1}(T_{-u_4}+jv)\\
AT_{-u_2} &= T_{u_2}\\
AT_{-u_3} &= T_{-u_1}\\
AT_{-u_4} &= T_{-u_4}
\end{aligned}$$

(3) $f(x) = x^2 + x + q$. Convention: $u_1 = v$, $u_2 = Av$, $u_3 = Av + v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=0}^{q-1}(T_{u_2}+jv) \cup \bigcup_{j=0}^{q-2}(T_{u_3}+jv)\\
AT_{u_2} &= T_{-u_3}\\
AT_{u_3} &= T_{-u_1}\\
AT_{-u_1} &= \bigcup_{j=0}^{q-1}(T_{-u_2}+jv) \cup \bigcup_{j=1}^{q-1}(T_{-u_3}+jv)\\
AT_{-u_2} &= T_{u_3}+(q-1)v\\
AT_{-u_3} &= T_{u_1}+(q-1)v
\end{aligned}$$

(4) $f(x) = x^2 - x + q$. Convention: $u_1 = v$, $u_2 = Av$, $u_3 = Av - v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=0}^{q-1}(T_{u_2} + jv) \cup \bigcup_{j=0}^{q-2}(T_{u_3} + jv) \\
AT_{u_2} &= T_{u_3} \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= \bigcup_{j=0}^{q-1}(T_{-u_2} + jv) \cup \bigcup_{j=1}^{q-1}(T_{-u_3} + jv) \\
AT_{-u_2} &= T_{-u_3} + (q-1)v \\
AT_{-u_3} &= T_{u_1} + (q-1)v
\end{aligned}$$

(5) $f(x) = x^2 + px + q$, $p \geq 2, 2p \leq q + 2$ (excluding $p = q = 2$). Convention:$u_1 = v$, $u_2 = Av + (p-1)v$, $u_3 = Av + pv$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=0}^{q-p}(T_{u_2} + jv) \cup \bigcup_{j=0}^{q-p-1}(T_{u_3} + jv) \\
AT_{u_2} &= \bigcup_{j=0}^{p-2}(T_{-u_2} + jv) \cup \bigcup_{j=0}^{p-1}(T_{-u_3} + jv) \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= \bigcup_{j=p-1}^{q-1}(T_{-u_2} + jv) \cup \bigcup_{j=p}^{q-1}(T_{-u_3} + jv) \\
AT_{-u_2} &= \bigcup_{j=q-p+1}^{q-1}(T_{u_2} + jv) \cup \bigcup_{j=q-p}^{q-1}(T_{u_3} + jv) \\
AT_{-u_3} &= T_{u_1} + (q-1)v
\end{aligned}$$

(6) $f(x) = x^2 - px + q$, $p \geq 2, 2p \leq q + 2$ (excluding $p = q = 2$). Convention: $u_1 = v$, $u_2 = Av - (p-1)v$, $u_3 = Av - pv$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=p-1}^{q-1}(T_{u_2} + jv) \cup \bigcup_{j=p}^{q-1}(T_{u_3} + jv) \\
AT_{u_2} &= \bigcup_{j=0}^{p}(T_{u_2} + jv) \cup \bigcup_{j=0}^{p-1}(T_{u_3} + jv) \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= \bigcup_{j=0}^{q-p}(T_{-u_2} + jv) \cup \bigcup_{j=0}^{q-p-1}(T_{-u_3} + jv) \\
AT_{-u_2} &= \bigcup_{j=q-p+1}^{q-1}(T_{-u_2} + jv) \cup \bigcup_{j=q-p}^{q-1}(T_{-u_3} + jv) \\
AT_{-u_3} &= T_{u_1} + (q-1)v
\end{aligned}$$

(7) $f(x) = x^2 + px - q,\ p \geq 1,\ 2p \leq q - 2$. Convention: $u_1 = v$, $u_2 = Av + pv$, $u_3 = Av + (p+1)v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=0}^{q-p-1} (T_{u_2} + jv) \cup \bigcup_{j=0}^{q-p-2} (T_{u_3} + jv) \\
AT_{u_2} &= T_{u_1} + (q-1)v \\
AT_{u_3} &= \bigcup_{j=q-p}^{q-1} (T_{u_2} + jv) \cup \bigcup_{j=q-p-1}^{q-1} (T_{u_3} + jv) \\
AT_{-u_1} &= \bigcup_{j=p}^{q-1} (T_{-u_2} + jv) \cup \bigcup_{j=p+1}^{q-1} (T_{-u_3} + jv) \\
AT_{-u_2} &= T_{u_1} \\
AT_{-u_3} &= \bigcup_{j=0}^{p-1} (T_{-u_2} + jv) \cup \bigcup_{j=0}^{p} (T_{-u_3} + jv)
\end{aligned}$$

(8) $f(x) = x^2 - px - q,\ p \geq 1,\ 2p \leq q - 2$. Convention: $u_1 = v$, $u_2 = Av - pv$, $u_3 = Av - (p+1)v$.

$$\begin{aligned}
AT_{u_1} &= \bigcup_{j=p}^{q-1} (T_{u_2} + jv) \cup \bigcup_{j=p+1}^{q-1} (T_{u_3} + jv) \\
AT_{u_2} &= T_{u_1} + (q-1)v \\
AT_{u_3} &= \bigcup_{j=q-p}^{q-1} (T_{-u_2} + jv) \cup \bigcup_{j=q-p-1}^{q-1} (T_{-u_3} + jv) \\
AT_{-u_1} &= \bigcup_{j=0}^{q-p-1} (T_{-u_2} + jv) \cup \bigcup_{j=0}^{q-p-2} (T_{-u_3} + jv) \\
AT_{-u_2} &= T_{-u_1} \\
AT_{-u_3} &= \bigcup_{j=0}^{p-1} (T_{u_2} + jv) \cup \bigcup_{j=0}^{p-2} (T_{u_3} + jv)
\end{aligned}$$

(9) $f(x) = x^2 + 2x + 2$. Convention: $u_1 = v$, $u_2 = Av + v$, $u_3 = Av + 2v$.

$$\begin{aligned}
AT_{u_1} &= T_{u_2} \\
AT_{u_2} &= T_{-u_2} \cup T_{-u_3} \cup (T_{-u_3} + v) \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= T_{-u_2} + v \\
AT_{-u_2} &= (T_{u_2} + v) \cup T_{u_3} \cup (T_{u_3} + v) \\
AT_{-u_3} &= T_{u_1} + v
\end{aligned}$$

(10) $f(x) = x^2 - 2x + 2$. Convention: $u_1 = v$, $u_2 = Av - v$, $u_3 = Av - 2v$.

$$\begin{aligned}
AT_{u_1} &= T_{u_2} + v \\
AT_{u_2} &= T_{u_2} \cup T_{u_3} \cup (T_{u_3} + v) \\
AT_{u_3} &= T_{-u_1} \\
AT_{-u_1} &= T_{-u_2} \\
AT_{-u_2} &= (T_{-u_2} + v) \cup T_{-u_3} \cup (T_{-u_3} + v) \\
AT_{-u_3} &= T_{u_1} + v
\end{aligned}$$

6. **Appendix C: Contact Matrices**

Let $f(x) = x^2 \pm px \pm q$ ($p \geq 0,\ q \geq 2$). The contact matrices (in table form) of disk-like tiles are classified by $f(x)$ and listed below.

	v	Av	$-v$	$-Av$	$Av - v$	$-Av - v$	$-Av + v$	$Av + v$
v	0	q	0	0	$q - 1$	0	0	$q - 1$
Av	0	0	1	0	0	0	0	0
$-v$	0	0	0	q	0	$q - 1$	$q - 1$	0
$-Av$	1	0	0	0	0	0	0	0
$Av - v$	0	0	0	0	0	1	0	0
$-Av - v$	0	0	0	0	0	0	1	0
$-Av + v$	0	0	0	0	0	0	0	1
$Av + v$	0	0	0	0	1	0	0	0

TABLE 4. $f(x) = x^2 + q$.

	v	Av	$-v$	$-Av$	$Av - v$	$-Av + v$	$Av + v$	$-Av - v$
v	0	q	0	0	$q - 1$	0	$q - 1$	0
Av	1	0	0	0	0	0	0	0
$-v$	0	0	0	q	0	$q - 1$	0	$q - 1$
$-Av$	0	0	1	0	0	0	0	0
$Av - v$	0	0	0	0	0	1	0	0
$-Av + v$	0	0	0	0	1	0	0	0
$Av + v$	0	0	0	0	0	0	1	0
$-Av - v$	0	0	0	0	0	0	0	1

TABLE 5. $f(x) = x^2 - q$.

	v	Av	$Av + v$	$-v$	$-Av$	$-Av - v$
v	0	q	$q - 1$	0	0	0
Av	0	0	0	0	0	1
$Av + v$	0	0	0	1	0	0
$-v$	0	0	0	0	q	$q - 1$
$-Av$	0	0	1	0	0	0
$-Av - v$	1	0	0	0	0	0

TABLE 6. $f(x) = x^2 + x + q$.

	v	Av	$Av-v$	$-v$	$-Av$	$-Av+v$
v	0	q	$q-1$	0	0	0
Av	0	0	1	0	0	0
$Av-v$	0	0	0	1	0	0
$-v$	0	0	0	0	q	$q-1$
$-Av$	0	0	0	0	0	1
$-Av+v$	1	0	0	0	0	0

Table 7. $f(x) = x^2 - x + q$.

	v	$Av+(p-1)v$	$Av+pv$	$-v$	$-Av-(p-1)v$	$-Av-pv$
v	0	$q-p+1$	$q-p$	0	0	0
$Av+(p-1)v$	0	0	0	0	$p-1$	p
$Av+pv$	0	0	0	1	0	0
$-v$	0	0	0	0	$q-p+1$	$q-p$
$-Av-(p-1)v$	0	$p-1$	p	0	0	0
$-Av-pv$	1	0	0	0	0	0

Table 8. $f(x) = x^2 + px + q,\ p \geq 2,\ 2p \leq q + 2$ (excluding $p = q = 2$).

	v	$Av-(p-1)v$	$Av-pv$	$-v$	$-Av+(p-1)v$	$-Av+pv$
v	0	$q-p+1$	$q-p$	0	0	0
$Av-(p-1)v$	0	$p-1$	p	0	0	0
$Av-pv$	0	0	0	1	0	0
$-v$	0	0	0	0	$q-p+1$	$q-p$
$-Av+(p-1)v$	0	0	0	0	$p-1$	p
$-Av+pv$	1	0	0	0	0	0

Table 9. $f(x) = x^2 - px + q,\ p \geq 2,\ 2p \leq q + 2$ (excluding $p = q = 2$).

	v	$Av+pv$	$Av+(p+1)v$	$-v$	$-Av-pv$	$-Av-(p+1)v$
v	0	$q-p$	$q-p-1$	0	0	0
$Av+pv$	1	0	0	0	0	0
$Av+(p+1)v$	0	p	$p+1$	0	0	0
$-v$	0	0	0	0	$q-p$	$q-p-1$
$-Av-pv$	0	0	0	1	0	0
$-Av-(p+1)v$	0	0	0	0	p	$p+1$

Table 10. $f(x) = x^2 + px - q,\ p \geq 1,\ 2p \leq q - 2$.

	v	$Av-pv$	$Av-(p+1)v$	$-v$	$-Av+pv$	$-Av+(p+1)v$
v	0	$q-p$	$q-p-1$	0	0	0
$Av-pv$	1	0	0	0	0	0
$Av-(p+1)v$	0	0	0	0	p	$p+1$
$-v$	0	0	0	0	$q-p$	$q-p-1$
$-Av+pv$	0	0	0	1	0	0
$-Av+(p+1)v$	0	p	$p+1$	0	0	0

Table 11. $f(x) = x^2 - px - q,\ p \geq 1,\ 2p \leq q - 2$.

	v	$Av+v$	$Av+2v$	$-v$	$-Av-v$	$-Av-2v$
v	0	1	0	0	0	0
$Av+v$	0	0	0	0	1	2
$Av+2v$	0	0	0	1	0	0
$-v$	0	0	0	0	1	0
$-Av-v$	0	1	2	0	0	0
$-Av-2v$	1	0	0	0	0	0

TABLE 12. $f(x) = x^2 + 2x + 2$.

	v	$Av-v$	$Av-2v$	$-v$	$-Av+v$	$-Av+2v$
v	0	1	0	0	0	0
$Av-v$	0	1	2	0	0	0
$Av-2v$	0	0	0	1	0	0
$-v$	0	0	0	0	1	0
$-Av+v$	0	0	0	0	1	2
$-Av+2v$	1	0	0	0	0	0

TABLE 13. $f(x) = x^2 - 2x + 2$.

REFERENCES

[AL] S. AKIYAMA AND B. LORIDANT, *Boundary parametrization of planar self-affine tiles with collinear digit set*, Sci. China Math. **53(9)** (2010), 2173-2194.

[AL2] S. AKIYAMA AND B. LORIDANT, *Boundary parametrization of self-affine tiles*, J. Math. Soc. Japan **63** no.2 (2011), 525-579.

[BG] C. BANDT AND G. GELBRICH, *Classification of self-affine lattice tilings*, J. London Math. Soc. **50** (1994), 581-593.

[BW] C. BANDT AND Y. WANG, *Disk-like self-affine tiles in* $\mathbb{R}^2$, Discrete Comput.Geom. **26** (2001), 591-601.

[DJN] D.-W. DENG, T. JIANG, AND S.-M. NGAI, *Structure of planar integral self-affine tilings*, Math. Nachr. **285** (2012), 447-475.

[DL] Q.R. DENG AND K.S. LAU, *Connectedness of a class of planar self-affine tiles*, J. Math. Anal. Appl. **380** (2011), 493-500.

[DKV] P. DUVAL, J. KEESLING, AND A. VINCE, *The Hausdorff dimension of the boundary of a self-similar tile*, J. London Math. Soc. **61** (2000), 748-760.

[Fa] K.J. FALCONER, *Fractal Geometry: Mathematical Foundations and Applications*, John Wiley & Sons, 2003.

[Fa2] K.J. FALCONER, *Techniques in fractal geometry*, John Wiley & Sons, 1997.

[Fi] R. FISCHER, *Sofic systems and graphs*, Monatsh. Math. **80** (1975), 179-186.

[Ga] F.R. GANTMACHER, *Matrix theory*, Volume II. Chelsea Publishing Company, New York, 1960.

[Gi] W.G. GILBERT, *Radix representataions of quadratic fields*, J. Math. Anal. Appl. **83** (1981), 264-274.

[HLR] X.G. HE, K.S. LAU, AND H. RAO, *Self-affine sets and graph-directed systems*, Constr. Approx. **19** (2003), 373-397.

[KL] I. KIRAT AND K.S. LAU, *On the connectedness of self-affine tiles*, J. London Math. Soc. **62** (2000), 291-304.

[LW1] J. C. LAGARIAS AND Y. WANG, *Self-affine tiles in* $\mathbb{R}^n$, Adv. Math. **121** (1996), 21-49.

[LW2] J. C. Lagarias and Y. Wang, *Integral Self-affine tiles in $\mathbb{R}^n$ I. Standard and non-standard digit sets*, J. London Math. Soc. **54** (1996), 161-179.

[LW3] J. C. Lagarias and Y. Wang, *Integral Self-affine tiles in $\mathbb{R}^n$ II. Lattice tilings*, J. Fourier Anal. Appl. **3** (1997), 84-102.

[LN] K.S. Lau and S.M. Ngai, *Dimension of the boundaries of self-similar sets*, Experiment. Math. **12** (2003), 13-26.

[LL] K.S. Leung and K.S. Lau, *Disk-likeness of planar self-affine tiles*, Trans. Amer. Math. Soc. **359** (2007), 3337-3355.

[LLu] K.S. Leung and J.J. Luo, *Connectedness of planar self-affine sets associated with non-consecutive collinear digit sets*, J. Math. Anal. Appl. (2012), In press.

[LM] D. Lind and B. Marcus, *An introduction to symbolic dynamics and coding*, Cambridge University Press, 1995.

[MW] R.D. Mauldin and S.C. Williams, *Hausdorff dimension in graph-directed constructions*, Trans. Amer. Math. Soc. **309** (1988), 811-829.

[MTT] W. Müller, J.M. Thuswaldner, and R.T. Tichy, *Fractal properties of number system*, Period. Math. Hunger. **42** (2001), 51-68.

[Se] E. Seneta, *Non-negative matrices and Markov chains*, Springer, New York, second edition, 1980.

[SW] R.S. Strichartz and Y. Wang, *Geometry of self-affine tiles I*, Indiana U. Math. J. **48** (1999), 1-24.

[We] B. Weiss, *Subshifts of finite type and sofic systems*, Monatsh. Math. **77** (1973), 462-474.

Department of Mathematics and Information Technology, The Hong Kong Institute of Education, Hong Kong
E-mail address: `ksleung@ied.edu.hk`

Department of Mathematics, The Chinese University of Hong Kong, Hong Kong
E-mail address: `jluo@math.cuhk.edu.hk`